

Power for linear models of longitudinal data with applications to Alzheimer's Disease Phase II study design

Michael C. Donohue, Steven D. Edland, Anthony C. Gamst
Division of Biostatistics and Bioinformatics
University of California, San Diego

January 11, 2013

1 Introduction

We will discuss power and sample size estimation for randomized placebo controlled studies in which the primary inference is based on the interaction of treatment and time in a linear mixed effects model (Laird and Ware 1982). We will demonstrate how the sample size formulas of Liu and Liang (1997) for marginal or generalized estimating equation (GEE) models (Zeger and Liang 1986) can be adapted for mixed effects models. Finally, using mixed effects model estimates based on data from the Alzheimer's Disease Neuroimaging Initiative (ADNI), we will give examples of sample size calculations for models with and without baseline covariates which may help explain heterogeneity in cognitive decline and improve power.

2 Power calculations

2.1 Exchangeable correlation and random intercept models

Suppose we wish to estimate the required sample size for inference regarding the interaction of treatment and time in a longitudinal, placebo controlled

study. Such calculations are relatively straightforward when the inference is based on a GEE model in which the correlation structure is assumed to be “exchangeable.” An exchangeable correlation structure specifies that all observations from within the same cluster, or repeated measures on the same subject, are equally correlated. This is exactly equivalent to a random effects model which includes a random intercept for each cluster of correlated observations. Sample sizes for study designs using these models can be calculated using a simple formula such as that in Diggle et al. 2002, page 29. The formula requires the number visits, the interval between visits, the estimated model variance (σ^2), the within subject correlation (ρ), and of course the usual sample size calculation inputs (power, significance level, and effect size).

To translate the formula of Diggle et al. 2002 to the random effects setting, let us first consider the details of the assumed error structure of the GEE framework. The GEE model assumes that the response for subject i at time t_{ij} , denoted Y_{ij} , is the group mean, dependent on time and treatment, plus an error term ε_{ij} . Or, borrowing notation from Diggle et al. 2002, for group A:

$$Y_{ij} = \beta_{0A} + \beta_{1A}t_{ij} + \varepsilon_{ij}, \quad i = 1, \dots, m; j = 1, \dots, n.$$

and similarly for Group B. The null hypothesis is $H_0 : d = \beta_{1A} - \beta_{1B} = 0$. Under an exchangeable correlation structure $\text{var}(Y_{ij}) = \text{var}(\varepsilon_{ij}) = \sigma^2$ and $\text{corr}(Y_{ij}, Y_{ik}) = \text{corr}(\varepsilon_{ij}, \varepsilon_{ik}) = \rho$, for all subjects, i , and time points j, k .

In the mixed effects framework we can assume a random intercept model which is equivalent to the GEE model with exchangeable correlation structure. In this case we believe $\varepsilon_{ij} = \alpha_i + \varepsilon_{ij}^*$, where α_i is the random intercept term shared by all observations and ε_{ij}^* are independent and identically distributed (iid) error terms. We see that $\text{var}(Y_{ij}) = \text{var}(\varepsilon_{ij}) = \text{var}(\alpha_i) + \text{var}(\varepsilon_{ij}^*)$ and $\text{corr}(Y_{ij}, Y_{ik}) = E[(\alpha_i + \varepsilon_{ij}^*)(\alpha_i + \varepsilon_{ik}^*)]/\sigma^2 = \text{var}(\alpha_i)/\sigma^2$. The variance of the random intercept, $\text{var}(\alpha_i)$, and the residual variance, $\text{var}(\varepsilon_{ij}^*)$, are easily obtainable from the output of mixed effects fitting software so that one might fit a random effects model to pilot data to educate a power calculation using the GEE formula of Diggle et al. 2002. Assuming equal numbers in the placebo and active groups, a common visit schedule for all subjects ($t_{ij} = t_{kj}$ for all i, j, k), and a random intercept model; the number of subjects per group is:

$$m = \frac{2(z_\alpha + z_Q)^2(\text{var}(\alpha_i) + \text{var}(\varepsilon_{ij}^*))^2(1 - \text{var}(\alpha_i)/\sigma^2)}{ns_x^2d^2}$$

where z_p is the p th standard normal quantile, Q is $1 - P$, P is the specified power, and $s_x^2 = n^{-1} \sum_j (t_j - \bar{x})^2$.

2.2 General correlation and random slope models

The random intercept model is not equipped to handle variations in the rate of change from subject to subject. In many diseases, such as Alzheimer's disease, the rate of improvement or decline will vary greatly within the treatment group, regardless of treatment. This variation can be modeled with a random slope term. That is, we assume:

$$Y_{ij} = \beta_{0A} + \beta_{1A}t_{ij} + \alpha_{0i} + \alpha_{1i}t_{ij} + \varepsilon_{ij}^*,$$

where we use ε_{ij}^* again to denote iid error and reserve ε_{ij} for possibly correlated error. If we derive the correlation structure of $\varepsilon_{ij} = \alpha_{0i} + \alpha_{1i}t_{ij} + \varepsilon_{ij}^*$, which is necessary in order to use GEE-based sample size formulas, we find that we no longer have an exchangeable correlation structure. In fact $\text{var}(Y_{ij}) = \text{var}(\varepsilon_{ij}) = \text{var}(\alpha_{0i}) + t_{ij}^2 \text{var}(\alpha_{1i}) + 2t_{ij} \text{cov}(\alpha_{0i}, \alpha_{1i}) + \text{var}(\varepsilon_{ij}^*)$ and $\text{cov}(Y_{ij}, Y_{ik}) = \text{cov}(\varepsilon_{ij}, \varepsilon_{ik}) = \text{var}(\alpha_{0i}) + t_{ij}t_{ik} \text{var}(\alpha_{1i}) + (t_{ij} + t_{ik}) \text{cov}(\alpha_{0i}, \alpha_{1i})$. For the common visit schedule case, the covariance matrix for the vector of correlated errors, $\boldsymbol{\varepsilon}_i = (\varepsilon_{i1}, \dots, \varepsilon_{in})'$, is of the form:

$$\Sigma = [(\text{var}(\alpha_0) + t_j t_k \text{var}(\alpha_1) + (t_j + t_k) \text{cov}(\alpha_0, \alpha_1))]_{jk} + \text{diag}(\text{var}(\varepsilon_j^*))$$

Armed with this specification of the covariance matrix one can use the sample size formula of Liu and Liang (1997) for linear GEE models (page 941). (Warning: The formula given on the bottom page 29 of Diggle et al. 2002 for general correlation matrices, R , is wrong).

The formula for linear models provided by Liu and Liang (1997) is useful for testing $H_0 : \boldsymbol{\psi} = \mathbf{0}$ for any linear model of the form:

$$Y_{ij} = \mathbf{x}_{ij}' \boldsymbol{\psi} + \mathbf{z}_{ij}' \boldsymbol{\lambda} + \varepsilon_{ij}$$

where $\boldsymbol{\varepsilon}_i \sim N(\mathbf{0}, \sigma^2 R)$ and the covariates for individual i , $\mathbf{x}_i = (\mathbf{x}_{i1}', \dots, \mathbf{x}_{in}')'_{n \times p}$ and $\mathbf{z}_i = (\mathbf{z}_{i1}', \dots, \mathbf{z}_{iq}')'_{n \times q}$, arise from a known discrete distribution. For our placebo controlled longitudinal study, the fully specified model is of the form:

$$Y_{ij} = \beta_0 + \beta_1 \{\text{Group}_i = A\} + \beta_2 t_{ij} + \beta_3 t_{ij} \{\text{Group}_i = A\}.$$

That is, the parameter of interest for the interaction of treatment and time is $\psi = \beta_3$ and nuisance parameter is $\boldsymbol{\lambda} = (\beta_0, \beta_1, \beta_2)'$. The covariates are

distributed as $\mathbf{x}_i = (t_1, \dots, t_n)'$ and $\mathbf{z}_j = [\mathbf{1} \mathbf{0} \mathbf{x}_i]_{n \times 3}$ with probability 1/2 (active group); and $\mathbf{x}_i = \mathbf{0}$ and $\mathbf{z}_j = [\mathbf{1} \mathbf{0} \mathbf{x}_i]_{n \times 3}$ with probability 1/2 (placebo group).

The Liu and Liang's formula for linear models can be coded In R as:

```
library(longpower)

liu.liang.linear.power

function (n = NULL, delta = NULL, u = NULL, v = NULL, sigma2 = 1,
  R = NULL, R.list = NULL, sig.level = 0.05, power = NULL,
  Pi = rep(1/length(u), length(u)), alternative = c("two.sided",
    "one.sided"))
{
  if (sum(sapply(list(n, delta, sigma2, power, sig.level),
    is.null)) != 1)
    stop("exactly one of 'n', 'sigma2', 'delta', 'power', and 'sig.level' must b
  if (!is.null(sig.level) && !is.numeric(sig.level) || any(0 >
    sig.level | sig.level > 1))
    stop("'sig.level' must be numeric in [0, 1]")
  alternative <- match.arg(alternative)
  if (sum(c(!is.null(R), !is.null(R.list))) != 1)
    stop("Exactly one of R or R.list must be specified.")
  if (sum(Pi) != 1)
    stop("Pi must sum to 1.")
  if (!is.null(R)) {
    R.list <- lapply(1:length(u), function(i) R)
  }
  Rinv <- lapply(1:length(R.list), function(i) {
    R <- R.list[[i]]
    if (is.null(dim(R)) & length(R) == 1 & length(u[[i]]) >
      1) {
      R <- matrix(R, length(u[[i]]), length(u[[i]])) +
        diag(1 - R, length(u[[i]]))
    }
    else if (is.null(dim(R)) & length(R) == 1 & length(u[[i]]) ==
      1) {
      R <- matrix(R, length(u[[i]]), length(u[[i]]))
    }
    return(solve(R))
  })
  n.body <- quote({
    Ipl <- 0
    for (i in 1:length(u)) Ipl <- Ipl + Pi[i] * t(u[[i]]) %**
      Rinv[[i]] %** v[[i]]
    Ipl <- Ipl/sigma2
  })
}
```

```

      Ill <- 0
      for (i in 1:length(u)) Ill <- Ill + Pi[i] * t(v[[i]]) **
        Rinv[[i]] ** v[[i]]
      Illinv <- solve(Ill/sigma2)
      Signal <- 0
      for (i in 1:length(u)) Signal <- Signal + Pi[i] * (t(u[[i]]) -
        Ipl ** Illinv ** t(v[[i]])) ** Rinv[[i]] ** (u[[i]] -
        v[[i]] ** Illinv ** t(Ipl))
      Signal <- Signal/sigma2
      ceiling((qnorm(1 - ifelse(alternative == "two.sided",
        sig.level/2, sig.level)) + qnorm(power))^2/(delta **
        Signal ** delta)[1, 1])
    })
  if (is.null(sig.level))
    sig.level <- uniroot(function(sig.level) eval(n.body) -
      n, c(1e-10, 1 - 1e-10))$root
  else if (is.null(power))
    power <- uniroot(function(power) eval(n.body) - n, c(0.001,
      1 - 1e-10))$root
  else if (is.null(delta))
    delta <- uniroot(function(delta) eval(n.body) - n, c(1e-10,
      1e+05))$root
  else if (is.null(sigma2))
    sigma2 <- uniroot(function(sigma2) eval(n.body) - n,
      c(1e-10, 1e+05))$root
  n <- eval(n.body)
  METHOD <- "Longitudinal linear model power calculation (Liu & Liang, 1997)"
  structure(list(n = n, delta = delta, sigma2 = sigma2, sig.level = sig.level,
    power = power, alternative = alternative, R = R, note = "n is *1/2* the total",
    method = METHOD), class = "power.longtest")
}
<environment: namespace:longpower>

```

The parameters include d , the effect size (possibly vector); u , the list of covariate vectors or matrices associated with the parameter of interest; v , the respective list of covariate vectors or matrices associated with the nuisance parameter; σ^2 , the error variance; R , the correlation structure; and P_i the proportion of covariates of each type (u , v , and P_i are expected to be the same length and sorted with respect to each other).

For example, we can reproduce the table exchangeable correlations on page 29 of Diggle et al for the case of $\mathbf{t} = (0, 2, 5)'$, $\alpha = 0.05$, power=0.80, and $d = 0.5$ as follows:

```

n = 3
t = c(0, 2, 5)

```

```

u = list(u1 = t, u2 = rep(0,n))
v = list(v1 = cbind(1,1,rep(0,n)),
        v2 = cbind(1,0,t))
rho = c(0.2, 0.5, 0.8)
sigma2 = c(100, 200, 300)
tab = outer(rho, sigma2,
            Vectorize(function(rho, sigma2){
              round(diggle.linear.power(
                d=0.5,
                t=t,
                sigma2=sigma2,
                R=rho,
                alternative="one.sided",
                power=0.80)$n)))
colnames(tab) = paste("sigma2 =", sigma2)
rownames(tab) = paste("rho =", rho)
tab

```

	sigma2 = 100	sigma2 = 200	sigma2 = 300
rho = 0.2	313	625	938
rho = 0.5	196	391	586
rho = 0.8	79	157	235

As a second example, consider an Alzheimer's disease trial in which assessments are taken every three months for 18 months (7 visits). We assume an smallest detectable effect size of 1.5 points on the cognitive portion of the Alzheimer's Disease Assessment Scale (ADAS-Cog). This is a 70 point scale with great variability among sick individuals. We assume the random intercept to have a variance of 55, the random slope to have a variance of 24, and a residual variance of 10. The correlation between random slope term and random intercept term is 0.8. We can estimate the necessary sample size by first generating the correlation structure. Since $\varepsilon = \text{var}(Y_{ij})$ is not constant over time in this model, we fix `sigma2=1` and set `R` equal to the covariance matrix for ε_i :

```

# var of random intercept
sig2.i = 55
# var of random slope
sig2.s = 24
# residual var
sig2.e = 10
# covariance of slope and intercep
cov.s.i <- 0.8*sqrt(sig2.i)*sqrt(sig2.s)
cov.t <- function(t1, t2, sig2.i, sig2.s, cov.s.i){

```

```

        sig2.i + t1*t2*sig2.s + (t1+t2)*cov.s.i
    }
t = seq(0,1.5,0.25)
n = length(t)
R = outer(t, t, function(x,y){cov.t(x,y, sig2.i, sig2.s, cov.s.i)})
R = R + diag(sig2.e, n, n)
u = list(u1 = t, u2 = rep(0,n))
v = list(v1 = cbind(1,1,rep(0,n)),
        v2 = cbind(1,0,t))
liu.liang.linear.power(d=1.5, u=u, v=v, R=R, sig.level=0.05, alternative="two.sided")

Longitudinal linear model power calculation (Liu & Liang, 1997)

        n = 208
        delta = 1.5
        sigma2 = 1
        sig.level = 0.05
        power = 0.8
        alternative = two.sided

```

NOTE: n is *1/2* the total sample size required.

R:

	[,1]	[,2]	[,3]	[,4]	[,5]	[,6]	[,7]
[1,]	65.00000	62.26636	69.53272	76.79908	84.06544	91.3318	98.59817
[2,]	62.26636	81.03272	79.79908	88.56544	97.33180	106.0982	114.86453
[3,]	69.53272	79.79908	100.06544	100.33180	110.59817	120.8645	131.13089
[4,]	76.79908	88.56544	100.33180	122.09817	123.86453	135.6309	147.39725
[5,]	84.06544	97.33180	110.59817	123.86453	147.13089	150.3972	163.66361
[6,]	91.33180	106.09817	120.86453	135.63089	150.39725	175.1636	179.92997
[7,]	98.59817	114.86453	131.13089	147.39725	163.66361	179.9300	206.19633

So the study would require about 208 subjects per arm to achieve 80% power, with a two-tailed $\alpha = 0.05$.

The simple formula provided in Diggle et al. 2002 suggests the required number of subjects can be found by $2(z_\alpha + 2Q)\xi/d^2$, where

$$\xi_{\text{WRONG}} = \begin{pmatrix} 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & \cdots & 1 \\ t_1 & \cdots & t_n \end{pmatrix} R^{-1} \begin{pmatrix} 1 & t_1 \\ \vdots & \vdots \\ 1 & t_n \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

Executing this for our Alzheimer's example, we get a sample size of:

```

x = (rbind(1,t) %*% solve(R) %*% cbind(1,t)) [2,2]
x*2*(qnorm(1-0.05/2) + qnorm(0.80))^2/1.5^2

```

[1] 0.3592744

which is clearly wrong. In fact, there is a typo in Diggle, et al (2002). The correct formula for ξ is:

$$\xi = \begin{pmatrix} 0 & 1 \end{pmatrix} \left[\begin{pmatrix} 1 & \cdots & 1 \\ t_1 & \cdots & t_2 \end{pmatrix} (\sigma^2 R)^{-1} \begin{pmatrix} 1 & t_1 \\ \vdots & \vdots \\ 1 & t_m \end{pmatrix} \right]^{-1} \begin{pmatrix} 0 \\ 1 \end{pmatrix}. \quad (1)$$

Applying the correct formula, we get

```
x = solve(rbind(1,t) %*% solve(R) %*% cbind(1,t)) [2,2]
x*2*(qnorm(1-0.05/2) + qnorm(0.80))^2/1.5^2
```

[1] 207.3101

Similary, using Liu and Liang 1997, we attempt to derive the correct closed form formula for this specific linear model. The required sample size per group is given as

$$m = \nu / (\psi_1' \tilde{\Sigma}_1 \psi_1)$$

where

$$\tilde{\Sigma}_1 = \sigma^{-2} \sum_{l=1}^m \pi_l (\mathbf{u}'_l - I_{\psi\lambda} I_{\lambda\lambda}^{-1} \mathbf{v}'_l) R^{-1} (\mathbf{u}'_l - \mathbf{v}_l I_{\lambda\lambda}^{-1} I'_{\psi\lambda}),$$

$$I_{\psi\lambda} = \sigma^{-2} \sum_{i=1}^m \pi_i \mathbf{u}'_i R^{-1} \mathbf{v}_i,$$

and

$$I_{\lambda\lambda} = \sigma^{-2} \sum_{i=1}^m \pi_i \mathbf{v}'_i R^{-1} \mathbf{v}_i.$$

Again, in our case the probability of each of the two covariate values is $\pi_1 = \pi_2 = 1/2$; and $\mathbf{u}_1 = (t_1, \dots, t_n)'$, $\mathbf{v}_1 = [\mathbf{1} \ \mathbf{0} \ \mathbf{x}_i]_{n \times 3}$, $\mathbf{u}_2 = \mathbf{0}$, and $\mathbf{v}_2 = [\mathbf{1} \ \mathbf{0} \ \mathbf{x}_i]_{n \times 3}$. We have

$$\begin{aligned} I_{\psi\lambda} &= \sigma^{-2} / 2 \mathbf{u}'_1 R^{-1} \mathbf{v}_1 \\ I_{\lambda\lambda} &= \sigma^{-2} / 2 [\mathbf{v}'_1 R^{-1} \mathbf{v}_1 + \mathbf{v}'_2 R^{-1} \mathbf{v}_2] = 1/2X \\ I_{\psi\lambda} I_{\lambda\lambda}^{-1} &= \mathbf{u}'_1 R^{-1} \mathbf{v}_1 X^{-1} \\ I_{\lambda\lambda}^{-1} I'_{\psi\lambda} &= X^{-1} \mathbf{v}'_1 R^{-1} \mathbf{u}_1 \end{aligned}$$

$$\begin{aligned}
\tilde{\Sigma}_1 &= \sigma^{-2}/2[(\mathbf{u}_1 - \mathbf{u}'_1 R^{-1} \mathbf{v}_1 X^{-1} \mathbf{v}'_1) R^{-1} (\mathbf{u}_1 - \mathbf{v}_1 X^{-1} \mathbf{v}'_1 R^{-1} \mathbf{u}_1) \\
&\quad + \mathbf{u}'_1 R^{-1} \mathbf{v}_1 X^{-1} \mathbf{v}'_2 R^{-1} \mathbf{v}_2 X^{-1} \mathbf{v}_1 R^{-1} \mathbf{u}_1 \\
&= \sigma^{-2}/2[\mathbf{u}_1 R^{-1} \mathbf{u} - \mathbf{u}'_1 R^{-1} \mathbf{v}_1 X^{-1} \mathbf{v}'_1 R^{-1} \mathbf{u}_1]
\end{aligned}$$

Applying this to our working example:

```

X = t(v[[1]])**solve(R)**v[[1]] +
    t(v[[2]])**solve(R)**v[[2]]
Sigma1 = ((t(u[[1]])**solve(R)**t(u[[1]])**solve(R)**v[[1]]**solve(X)**t(v[[1]]))
(qnorm(1-0.05/2) + qnorm(0.80))^2/(Sigma1*(1.5)^2)

[1, 1]
[1, ] 207.3101

```

3 References

- Diggle PJ, Heagerty PJ, Liang K-, Zeger SL. Analysis of longitudinal data. Second Edition. Oxford Statistical Science Series. 2002.
- Laird, N.M. and Ware, J.H. (1982) Random-Effects Models for Longitudinal Data, Biometrics, 38, 963-974.
- Liu G, Liang KY. Sample size calculations for studies with correlated observations. Biometrics. 1997;53:937-47.
- Zeger, S.L. and Liang, K.Y. (1986) Longitudinal data analysis for discrete and continuous outcomes. Biometrics, 42 121–130.